

Effect of Particle Size and Extraction Time on Total Flavonoid Level of *Artemisia vulgaris* Ethanol Extract

^{1*}Kholif Sholehah Indra Kurniasih, ²Mitsalina Fildzah Arifah, ³Eka Ayu Setiyawati, ⁴Syafira Atika Rahmah, ⁵Lucia Widianingrum, ⁶Eva Kholifah

¹⁻²Departement of Pharmacy, Faculty of Health, Jenderal Achmad Yani Yogyakarta University

³⁻⁵Bachelor, Departement of Pharmacy, Faculty of Health, Jenderal Achmad Yani Yogyakarta University

⁶Department of Pharmaceutical Chemistry, Pharmacy Study Program, Salsabila College of Health Sciences, Serang

¹kholifsholehahindra@gmail.com*; ²mitsalina.fildzah.arifah@gmail.com; ³ekaayusetiyawati@gmail.com; ⁴syafiraatikarahmah@gmail.com;

⁵luciawidhianngrum02@gmail.com; ⁶evakholifah.apt@gmail.com

*corresponding author

ARTICLE INFO

Article history

Received: November 23rd 2024

Revised: December 29th 2024

Accepted: December 31st 2024

Keywords

Artemisia vulgaris

Total Flavonoids Content

Particle Sizes

Extraction Time

ABSTRACT

Background: *Artemisia vulgaris*, also referred to as Daun Baru Cina, is a herbaceous species within the Asteraceae family. *Artemisia vulgaris* is a traditional remedy for numerous ailments, exhibiting anti-inflammatory and antioxidant properties due to its secondary metabolites, specifically flavonoids. The extraction method is essential for extracting the desired chemical from natural sources intended for therapeutic use. The extraction process is affected by several elements, including the particle size of simplicia powder and the duration of extraction. Objective: This study aims to test the effect of differences in particle size of powdered simplicia and extraction time of *Artemisia vulgaris* leaves on total flavonoid levels, using 70% ethanol solvent. Researchers used a quantitative approach using a descriptive research design. In this study, a qualitative test was carried out in the form of phytochemical screening followed by a quantitative test to determine the total flavonoid levels of *Artemisia vulgaris* leaves. Various particle sizes (40 mesh and 80 mesh) and extraction times (12 hours and 36 hours) were used in this study. Using ultraviolet-visible spectrophotometry, the Total Flavonoid Level (TFC) of the ethanol extract of *Artemisia vulgaris* leaves was determined. Results: The TFC value obtained for the optimal particle size was 40 mesh, with an extraction time of 12 hours, which was 72.073 ± 1.126 mgQE / g extract. While 80 mesh, with an extraction time of 36 hours, gave a TFC value of 70.169 ± 0.480 mgQE / g extract. In addition, 70% ethanol extract of *Artemisia vulgaris* leaves contains secondary metabolites of phenolic alkaloids, tannins, saponins, and steroids. Conclusion: The study concludes that the quantity of simplicia powder particles and the extraction duration influence the total flavonoid concentrations in *Artemisia vulgaris*.

This is an open access article under the [CC-BY-SA](#) license.

How to Cite: Kurniasih, K.S.I., Arifah, M.F., Setiyawati, E.A., Rahmah, S.A., Widianingrum, L., & Kholifah, E. (2024). Effect of Particle Size and Extraction Time on Total Flavonoid Level of *Artemisia vulgaris* Ethanol Extract. *Journal of Biotechnology and Natural Science*, 4(2): 72-80.

1. Introduction

Products originating from nature can be obtained from various sources, such as plants, animals, and microorganisms, which contain a large number of bioactive compounds (Abiri et al., 2018). This is



because bioactive compounds in plants can treat many diseases. In particular, plants have the potential to be utilized as therapies for a variety of diseases due to the presence of numerous bioactive substances within them. According to research by Rivai et al., (2020), the term "traditional medicine" is widely used to refer to treatments that result from plants. According to Judzentiene & Garjonyte, (2016), an excellent possibility exists for medicinal plants to be utilized as alternative medications due to the fact that they are non-toxic and have a limited number of adverse effects. According to Anibogwu et al., (2021), the variety of *Artemisia vulgaris* is a type of *Artemisia* species that is extensively distributed over the continents of Africa, North and South America, Asia, and Europe (Anibogwu et al., 2021). *Artemisia vulgaris* has been proven to show effects as an antioxidant (Budiana et al., 2017a; Febrina et al., 2017), hypolipidemic (Khan & Abedulla Khan, 2015), hepatoprotective and cytotoxic (Baykan Erel et al., 2011), antibacterial (Puspita Sari et al., 2021), estrogenic, antispasmodic, and analgesic (Obistioiu et al., 2014; Pandey et al., 2021). The *Artemisia vulgaris* type is less explored because it contains lower levels of bioactive compounds compared to other types of *Artemisia* (Ekiert et al., 2020). However, the use of natural ingredients can be used as a preliminary study and developed into natural medicinal preparations.

Artemisia vulgaris contains bioactive compounds such as flavonoids, phenolics (Dhifallah, 2022), essential oils (Bora & Sharma, 2011; Judzentiene & Garjonyte, 2016; Obistioiu et al., 2014), sterols, carotenoids and coumarins (Trifan et al., 2022). Flavonoids are one of the most studied types of secondary metabolites and provide various biological activities such as anti-inflammatory, anticancer, and antioxidant (Reungoat et al., 2021). The content of secondary metabolites such as flavonoids obtained can be influenced by several factors, such as extraction method, type of solvent, particle size, and length of extraction time (Syamsul et al., 2019). It has been asserted in the research that was reported by Malau et al., (2021); Tiwari et al., (2023) and Xiao et al., (2014) that the flavonoid content in *Artemisia vulgaris* and *Artemisia lactiflora* can be affected by the concentration of ethanol dissolution and the extraction time. There is still no research on how the particle size and maceration time of a 70% ethanol extract of *Artemisia vulgaris* affect the total amount of flavonoids. This research was conducted to evaluate the effect of variations in the particle size of simplicia powder and the maceration time of a 70% ethanol extract of *Artemisia vulgaris* on the total flavonoid content as a candidate for traditional medicine.

2. Methods

2.1 Preparation of simplicia

Artemisia vulgaris L. leaves were obtained from CV. Merapi Farma Herbal, Sleman, Yogyakarta. Dried *Artemisia vulgaris* leaves were cut into small pieces, crushed, and sieved at 40 and 80 mesh to ensure they were passed the 40 mesh sieve but not the 80 mesh sieve, but also produced powder that was passed the 80 mesh sieve.

2.2 Preparation of extracts

The sample powder was sieved and found to weigh 200 g. After undergoing the particle size variation treatment, it was extracted with 500 mL of 70% ethanol solvent at a ratio of 1:2.5. The sample powder was macerated at 25 °C for 12 and 36 hours before being filtered using a Buchner funnel. The filtrate obtained from the maceration results is evaporated until a thick extract is achieved. The yield % is determined by weighing the thick extract and applying the formula (Rahmawati et al., 2022).

$$\% \text{ Yield} = \frac{\text{Extract weight (g)}}{\text{Sample powder weight (g)}} \times 100\% \quad (1)$$

2.3 Phytochemical screening

Secondary metabolite compounds can be identified qualitatively, using the phytochemical screening method. Flavonoids, phenolics, alkaloids, tannins, saponins, and steroids are among the secondary metabolite compounds often found in plants.

2.3.1 Flavonoids

2 mL of 70% ethanol extract of *Artemisia vulgaris* leaves are heated for 5 minutes; 0.2 g magnesium and 5 drops of concentrated H₂SO₄ are added. A positive result for flavonoids is obtained when the color changes from yellow to red (Leswara et al., 2024).

2.3.2 Phenolic

A total of 2 mL of 70% ethanol extract from *Artemisia vulgaris* leaves was reacted with 2 drops of 5% FeCl₃. The color change should be observed; if a blackish-green color is formed, the presence of phenolic compounds in the extract is indicated. This reaction is characteristic of phenolic compounds, known for their antioxidant properties and potential health benefits (Leswara et al., 2024).

2.3.3 Alkaloids

5 mL of 70% ethanol extract of *Artemisia vulgaris* leaves is taken, 5 mL of 2 N HCl is added, then it is divided into 2 in a test tube. 3 drops of Dragendoff's reagent are added to the first tube, then 3 drops of Mayer's reagent are added to the second tube, and 3 drops of Wagner's reagent are added to the third tube. If an orange precipitate is formed in the first test tube, in the second test tube a white to yellow precipitate is formed, and in the third test tube a brown precipitate is formed, this indicates a positive alkaloid (Leswara et al., 2024).

2.3.4 Tannin

According to Ikalinus et al., (2015), the presence of a white precipitate is indicated by adding 1% gelatin solution after 2 mL of 70% ethanol *Artemisia vulgaris* extract reacts with 10 mL of 10% NaCl solution.

2.3.5 Saponins

According to Munadi & Arifin, (2022), a positive saponin result is indicated when the foam is occurred in the solution for at least 30 seconds after combining 10 mL of distilled water with 2 mL of 70% ethanol extracted from *Artemisia vulgaris* and shaking for approximately 30 seconds.

2.3.6 Steroids

One gram of extract was weighed, 2 mL of chloroform was added, and the mixture was agitated. The resulting filtrate was mixed with two drops of chloroform and one drop of concentrated H₂SO₄. Triterpenoid molecules are indicated by a red or purple color shift, while steroid compounds are indicated by a blue or green color change (Sari et al., 2021).

2.4 Determination of Total Flavonoid Content (TFC)

The total flavonoid extract levels in each treatment were found using the Chang method with UV-Vis spectrophotometry and quercetin as a standard. Subsequently, the results obtained are expressed in g QE/mg (Budiana et al., 2017b; Chang Chia-Chi et al., 2002).

2.4.1 Preparation of quercetin standard solution (1000 ppm)

After 50.0 mg of quercetin is measured, it is dissolved in ethanol p.a to the limit line in a 50.0 mL volumetric flask. The mixture is shaken until it is homogenous.

2.4.2 Determination of the optimum wavelength of quercetin

The maximum wavelength is determined by using a standard concentration of quercetin of 60 ppm, 1 mL is taken and placed in a 10 mL measuring flask. In the measuring flask, 1.0 mL of 10% AlCl₃ and 5% CH₃COOH were added to the limit line. The absorbance at λ 350-500 nm is then read (Nisaa et al., 2023).

2.4.3 Determination of the operating time (OT) of quercetin

A standard solution of 60 ppm was created from the standard 1000 ppm of quercetin. 1 mL of 60 ppm standard solution was placed in a 10 mL measuring flask, and then topped off with 1.0 mL of 10% aluminum chloride (AlCl₃) and 5% acetic acid (CH₃COOH). The absorption between 0 and 60 minutes was then measured at the highest wavelength achieved (Nisaa et al., 2023).

2.4.4 Standard curve for quercetin

A six concentration series: 30, 45, 60, 75, 90, and 105 ppm in ethanol p.a was created using a 1000 ppm quercetin standard solution. Each solution was added, 1 mL of 10% AlCl₃ and 5% CH₃COOH, to a 10.0 mL volumetric flask and filled. The absorbance at quercetin's optimal λ was measured after incubation with OT.

2.4.5 Determination of total flavonoid content of *Artemisia vulgaris* leaf extract

A stock solution of 1000 ppm extract was prepared for each treatment by weighing 100.0 mg of extract in a 10.0 mL measuring flask and adding ethanol solvent p.a. After that, 1 mL was transferred to a 10-mL measuring flask and the limit line was filled with 1.0 mL of 10% AlCl₃ and 5% CH₃COOH. Incubation was done during OT and absorbance was measured at the appropriate wavelength (Nisaa et al., 2023). Three replications of the extraction were performed, and Total Flavonoid Content (TFC) was calculated using the Chang method formula.

$$\text{TFC} = \frac{C \times V \times Fp}{W} \times 100\% \quad (2)$$

3. Results and Discussion

3.1 Sample preparation

Artemisia vulgaris L. leaves were obtained from CV. Merapi Farma Herbal, Sleman, Yogyakarta had previously determined to ensure that the leaves taken were *Artemisia vulgaris* L leaves. The determination was carried out at Merapi Farma Herbal Yogyakarta with the number SK - SMPL / MFH / VII / 2024. Based on the results of the determination, it was found that the sample used is Baru Cina leaves with the Latin name *Artemisia vulgaris*. Fresh leaves were dried using direct sunlight and covered with black cloth for five days. After that, the water content is checked using Moisture Balance. If the air content exceeds 10%, proceed with drying in an oven set to 40°C to obtain a moisture content of 5.8%. In this study, the water content of *Artemisia vulgaris* leaf simplicia was 5.80%, which meets the Ministry of Health's guideline for a water content of less than 10%. The dry material is ground into powder and sieved through mesh numbers 40 and 80. The goal of powdered simplicia is to increase the contact surface area between the sample and the solvent, speeding up the extraction process by reducing the solute diffusion distance. The sieving technique ensures that the size of the simplicia particles employed in the extraction process is consistent (Asworo & Widwiastuti, 2023). A finer simplicia particle size can improve extraction efficiency, but if the particles are too small, the likelihood of contaminants in the extract increases (Illing et al., 2023). In this study, the maceration method was used for extraction. The maceration procedure creates a pressure difference between the interior and outside of the simplicia cell, causing the cell wall and membrane to rupture. This allows the active chemicals in the cytoplasm to be removed and dissolved in the solvent utilized (Handoyo D.L.Y, 2020).

70% ethanol solvent was used because 70% ethanol has good solubility for many bioactive compounds, including flavonoids. Ethanol is a polar solvent. Therefore, it can dissolve polyphenolic substances like flavonoids, known to dissolve in polar solvents. A 70% ethanol solvent was used to sift the dried leaves through 40 and 80 mesh sieves, with each sieve number macerating for 12 and 36 hours. The yield percentage value was calculated by comparing the thick extract results to the beginning weight of the simplicia, as shown in Table 1.

Table 1. Results of extraction of *Artemisia vulgaris* leaves

Sieve Number	Maceration Time (hours)	%Yield
40	12	1.580
40	36	1.190
80	12	2.975
80	36	1.895

Table 1 shows that the average yield value of *Artemisia vulgaris* leaf extract increases with increasing particle size and maceration time. This is consistent with the hypothesis that the smaller the particle size, the higher the surface area available for interaction with the solvent. The extraction procedure involves transferring active chemicals from the cell to the solvent. greater active chemicals can be recovered from smaller particles because the greater surface area is exposed to the solvent (Gil-Martín et al., 2022). However, the yield fell at a particle size of 80 mesh and a maceration time of 36 hours. This can be produced by over-saturation during extraction. After 36 hours of extraction, the solvent may approach a saturation point (over-saturation) with the active molecule, which means that the solvent can no longer dissolve new compounds. When the solvent reaches saturation, extending the extraction time will not increase the amount of substance removed and may reduce extraction efficiency, lowering yield. Smaller particle size can increase extraction, however excessive extraction time can reduce extraction efficiency.

3.2 Phytochemical screening

The phytochemical screening of a 70% ethanol extract of *Artemisia vulgaris* in each treatment yielded positive results containing flavonoids, phenolics, alkaloids, tannins, saponins, and steroids, as illustrated in Table 2.

Table 2. Phytochemical screening of 70% ethanol extract of *Artemisia vulgaris*

Compound	Reagent	Results	Description
Flavonoids	H ₂ SO ₄ p	yellow brown color	+
Phenolic	FeCl ₃ 5%	blackish green color	+
Alkaloids	Mayer	yellowish white precipitate	+
	Dragendorff	red precipitate	+
	Wagner	Brown precipitate	+
Tannin	NaCl + Gelatin	yellowish precipitate	+
Saponin	Aquadest + HCl 1 N	foam	+
Steroid	Chloroform + H ₂ SO ₄	red ring	+

Note: (+) positive containing secondary metabolites

An interaction between concentrated sulfuric acid (H₂SO₄) and flavonoids is possible due to their hydroxyl group (-OH). Depending on the kind of flavonoid, adding sulfuric acid to flavonoid molecules causes a chromogenic reaction that can result in a yellow, purple, or red color. Sulfuric acid is involved in the dehydration of flavonoids and forms a compound with them, responsible for their color. For example, sulfuric acid can degrade flavonoids by releasing air molecules, which can alter their structure into more stable compounds and give them certain colors.

The reaction between tannin, NaCl, and gelatin produces precipitation resulting from the formation of a complex between tannin and protein in gelatin. The reaction between steroids, sulfuric acid (H₂SO₄), and chloroform can be used in a qualitative test to detect the presence of steroids in samples. This reaction is known as the Salkowski test. Sulfuric acid can oxidize sterols in the sample to form complex compounds, causing a change in the red ring in the chloroform layer.

3.3 Determination of the maximum wavelength and OT of quercetin

In this study, the maximum wavelength was determined in the range of 0–60 minutes, and a maximum wavelength of 414 nm was obtained. The optimum wavelength is determined when a complex between quercetin – AlCl₃ occurs, which provides optimum absorption. Measurement at the maximum wavelength allows for the largest change in absorbance for each unit level so as to minimize measurement errors and increase the accuracy of results when replication or repeated measurements are carried out (Suharyanto & Prima D.A.N, 2020). Meanwhile, the operating time (OT) of quercetin was determined at a concentration of 60 ppm using the optimum wavelength obtained, namely 414 nm. The measurement data show that the operating time (OT) starts from 42 to 47 minutes, and an absorbance value of 0.397 is obtained. Similar results were also found in research by Pujiastuti & Ma'rifah (2022), where the operating time for quercetin was 40 minutes. Determining the operating time aims to determine the time at which absorbance reaches maximum stability, which indicates that the reaction between compounds has reached an equilibrium point. This is important to minimize errors in measurement (Suharyanto & Prima D.A.N, 2020).

Several concentration series of 1000 ppm quercetin stock curves created a quercetin standard curve, including 30, 45, 60, 75, 90, and 105 ppm. Next, in each concentration series, the absorbance was measured at a wavelength of 414 nm, and measurements were carried out in the OT range of 42–47 minutes. The absorbance obtained from each concentration series can be seen in Table 3.

Table 3. Standard curve series of quercetin

Concentration (ppm)	Absorbance			Absorbance Average
	1	2	3	
30	0.189	0.182	0.185	0.185
45	0.263	0.265	0.263	0.264
60	0.339	0.338	0.347	0.341
75	0.469	0.452	0.452	0.458
90	0.571	0.538	0.550	0.553
105	0.656	0.670	0.667	0.664

From the results of absorbance data analysis at various concentrations, a linear regression equation $y = 0.0064x - 0.0236$ was obtained with a correlation coefficient (r) of 0.9973. An r value that is greater than the r table (0.8783) indicates a linear relationship between concentration and absorbance. The higher the standard concentration of quercetin, the higher the absorption value. The choice of concentration is based on the Lambert-Beer law which states that the absorption requirement is 0.2 to 0.8. The resulting standard curve will be a straight line if the Lambert-Beer

law is fulfilled (Grace et al., 2015). This equation is valid and can be used to determine the ethanol 70% total flavonoid content of *Artemisia vulgaris* leaf extract in each treatment. The calibration curve graph is shown in Figure 1.




Figure 1. Calibration curve of quercetin

3.4 Determination of flavonoid content of 70% ethanol extract of *Artemisia vulgaris* leaves

The total flavonoid content of a 70% ethanol extract of *Artemisia vulgaris* leaves was quantified using the UV-Vis spectrophotometric technique at a wavelength of 414 nm. The samples were then incubated for 42–47 minutes, depending on the results. The total flavonoid concentrations were determined using the derived calibration curve. The test results indicated that the average total flavonoid content for sieve number 40 with a maceration duration of 12 hours was $\bar{x} \pm \text{LE} = 72.073 \pm 1.126$ mgQE/g extract, for sieve number 40 with a maceration duration of 36 hours was $\bar{x} \pm \text{LE} = 65.645 \pm 0.229$ mgQE/g extract, for sieve number 80 with a maceration duration of 12 hours was $\bar{x} \pm \text{LE} = 69.70 \pm 0.278$ mgQE/g extract, and for sieve number 80 with a maceration duration of 36 hours was $\bar{x} \pm \text{LE} = 70.169 \pm 0.480$ mgQE/g extract. The assay results are presented in Table 4.

Table 4. Determination of total flavonoid content

Particle Size (mesh)	Maceration Time (hours)	
	12 (mgQE/g extract)	36 (mgQE/g extract)
40	72.073 ± 1.126	65.645 ± 0.229
80	69.70 ± 0.278	70.169 ± 0.480

The average amount of flavonoids in *Artemisia vulgaris* leaves was highest in the 40 mesh particle size treatment, which was macerated for 12 hours and had a value of 72.073 ± 1.126 mgQE/g extract (Table 4). A larger particle size can provide good permeability for solvents, which allows flavonoid compounds to dissolve more efficiently without the risk of dissolved impurities. Finer particles, specifically those with an 80 mesh particle size, offer a greater surface area. However, they often dissolve impurity compounds alongside active compounds, potentially lowering the quality of the extract. In addition, in very fine particles, apart from flavonoids, other compounds such as salts or proteins can also dissolve, which can reduce the levels of pure flavonoids that will be obtained (Hijaz et al., 2020).

4. Conclusion

The conclusions of the study indicate that secondary metabolites, including flavonoids, phenolics, tannins, alkaloids, saponins, and steroids, are present in 70% ethanol extract of *Artemisia vulgaris* leaves. The most effective method for extracting secondary metabolites from *Artemisia vulgaris* leaves is the treatment that involves a particle size of 40 mesh and a maceration duration of 12 hours with a total flavonoid content of $\bar{x} \pm \text{LE} = 72.073 \pm 1.126$ mgQE/g extract

REFERENCES

- Abiri, R., Silva, A. L. M., de Mesquita, L. S. S., de Mesquita, J. W. C., Atabaki, N., de Almeida, E. B., Shaharuddin, N. A., & Malik, S. (2018). Towards a better understanding of *Artemisia vulgaris*: Botany, phytochemistry, pharmacological and biotechnological potential. *Food Research International*, 109, 403–415. <https://doi.org/10.1016/J.FOODRES.2018.03.072>.
- Anibogwu, R., De Jesus, K., Pradhan, S., Pashikanti, S., Mateen, S., & Sharma, K. (2021). Extraction, isolation and characterization of bioactive compounds from artemisia and their biological significance: A review. In *Molecules* (Vol. 26, Issue 22). MDPI. <https://doi.org/10.3390/molecules26226995>.
- Asworo, R. Y., & Widwastuti, H. (2023). Pengaruh Ukuran Serbuk Siplisia dan Waktu Maserasi terhadap Aktivitas Antioksidan Ekstrak Kulit Sirsak. *Indonesian Journal of Pharmaceutical Education*, 3(2). <https://doi.org/10.37311/ijpe.v3i2.19906>.
- Baykan Erel, Ş., Gökhan Şenol, S., Köse, F. A., & Ballar, P. (2011). In Vitro Cytotoxic Properties Of Six *Artemisia* L. Species. In *Şüra BAYKAN EREL, Serdar Gökhan ŞENOL* (Vol. 8, Issue 3).
- Bora, K. S., & Sharma, A. (2011). The Genus *Artemisia*: A Comprehensive Review. *Pharmaceutical Biology*, 49(1), 101–109. <https://doi.org/10.3109/13880209.2010.497815>.
- Budiana, W., Suhardiman, A., Roni, A., Sumarah, I., & Nara, T. E. (2017a). Aktivitas antioksidan ekstrak daun tiga genus *Artemisia* sp dengan metode DPPH serta penetapan kadar total flavonoid, fenol dan karotenoid. *Kartika Jurnal Ilmiah Farmasi*, 5(2), 38. <https://doi.org/10.26874/kjif.v5i2.106>.
- Budiana, W., Suhardiman, A., Roni, A., Sumarah, I., & Nara, T. E. (2017b). Aktivitas antioksidan ekstrak daun tiga genus *Artemisia* sp dengan metode DPPH serta penetapan kadar total flavonoid, fenol dan karotenoid. *Kartika Jurnal Ilmiah Farmasi*, 5(2), 38. <https://doi.org/10.26874/kjif.v5i2.106>.
- Chang Chia-Chi, Yang Ming-Hua, Wen Hwei-Mei, & Chern Jiing-Chuang. (2002). Estimation of Total Flavonoid Content in Propolis by Two Complementary Colorimetric Methods. *Journal of Food and Drug Analysis*, 10(3), 178–182.
- Dhifallah, A., S. H., O. A., S. H., A. D., & R. H. (2022). Comparative Study of Phenolic Compounds and Antiradical Activities of Four Extracts of Tunisian *Artemisia herba Alba*. *Pharm. Chem. J.*, 56, 226–232.
- Ekiert, H., Pajor, J., Klin, P., Rzepiela, A., Slesak, H., & Szopa, A. (2020). Significance of *artemisia vulgaris* L. (*Common Mugwort*) in the history of medicine and its possible contemporary applications substantiated by phytochemical and pharmacological studies. In *Molecules* (Vol. 25, Issue 19). MDPI AG. <https://doi.org/10.3390/molecules25194415>.
- Febrina, L., Riris, I. D., & Silaban, S. (2017). Uji aktivitas antibakteri terhadap *Escherichia coli* dan antioksidan dari ekstrak air tumbuhan binara (*Artemisia vulgaris* L.). *Jurnal Pendidikan Kimia*, 9(2), 311–317. <https://doi.org/10.24114/jpkim.v9i2.7621>.
- Gil-Martín, E., Forbes-Hernández, T., Romero, A., Cianciosi, D., Giampieri, F., & Battino, M. (2022). Influence of the extraction method on the recovery of bioactive phenolic compounds from food industry by-products. In *Food Chemistry* (Vol. 378). Elsevier Ltd. <https://doi.org/10.1016/j.foodchem.2021.131918>.
- Grace, K., Davenport, F., Hanson, H., Funk, C., & Shukla, S. (2015). Linking climate change and health outcomes: Examining the relationship between temperature, precipitation and birth weight in Africa. *Global Environmental Change*, 35, 125–137. <https://doi.org/10.1016/J.GLOENVCHA.2015.06.010>.
- Handoyo D.L.Y. (2020). Pengaruh Lama Waktu Maserasi (Perendaman) Terhadap Kekentalan Ekstrak Daun Sirih (*Piper betle*) The Influence of Maseration Time (Immeration) on The Vocity of Birthleaf Extract (*Piper betle*). *Jurnal Farmasi Tinctura*, 2(1), 34–41. <https://doi.org/https://doi.org/10.35316/tinctura.v2i1.1546>.

- Hijaz, F., Al-Rimawi, F., Manthey, J., & Killiny, N. (2020). Phenolics, flavonoids and antioxidant capacities in *Citrus* species with different degree of tolerance to Huanglongbing. *Plant Signaling & Behavior*, *15*, 1752447. <https://doi.org/10.1080/15592324.2020.1752447>.
- Ikalinus, R., Widyastuti, S., & Eka Setiasih, N. (2015). Skrining Fitokimia Ekstrak Etanol Kulit Batang Kelor (*Moringa oleifera*). *Indonesia Medicus Veterinus*, *4*(1), 71–79.
- Illing, I., Iman, F., & Sukarti. (2023). Analisis Kadar Flavonoid Ekstrak Rumput KNOP (*Hyptis capitata* jacq) dengan Metode Spektrofotometri UV-VIS. *Cokroaminoto Journal of Chemical Science*, *5*(1), 20–24.
- Judzentiene, A., & Garjonyte, R. (2016). Compositional Variability and Toxic Activity of Mugwort (*Artemisia vulgaris*) Essential Oils.
- Khan, K. A., & Abedulla Khan, K. (2015). A preclinical antihyperlipidemic evaluation of *Artemisia vulgaris* root in diet induced hyperlipidemic animal model. *5*, 110. <https://doi.org/10.7439/ijpr>.
- Leswara, D. F., I, K. S., & Kurniasih. (2024). The Effect of Maceration Duration on The Total Flavonoids Content of *Kaempferia parviflora* wall. ex baker Ethanol Extract. *MEDIA ILMU KESEHATAN*, *13*(1). <https://doi.org/10.30989/mik.v13i1.1300>.
- Malau, A. G., Widyasanti, A., & Putri, S. H. (2021). Optimization of Ultrasonic Assisted Extraction Process on Antioxidant Activity of Honje Fruit Extract (*Etilingera elatior*) using Surface Response Method. *Jurnal Kimia Valensi*, *7*(2), 118–128. <https://doi.org/10.15408/jkv.v7i2.21396>.
- Munadi, R., & Arifin, L. (2022). Identifikasi Senyawa Metabolit Sekunder dan Uji Aktivitas Antioksidan Ekstrak Daun Jahe Putih (*Zingiber officinale* Rosc. var. officinarum). *Spin*, *4*(2), 163–174. <https://doi.org/10.20414/spin.v4i2.5420>.
- Nisaa, N. R. K., Malik, Abd., & Handayani, V. (2023). Analisis Kadar Total Flavonoid Ekstrak Etanol Kulit Pisang Cavendish (*Musa paradisiaca* var. Sapiantum) Menggunakan Metode Spektrofotometri Uv-Vis. *Jurnal Sains Dan Kesehatan*, *5*(2), 212–217. <https://doi.org/10.25026/jsk.v5i2.1810>.
- Obistoiu, D., Cristina, R. T., Schmerold, I., Chizzola, R., Stolze, K., Nichita, I., & Chiurciu, V. (2014). Chemical Characterization by GC-MS and in vitro activity against *Candida albicans* of volatile fractions prepared from *Artemisia dracunculus*, *Artemisia abrotanum*, *Artemisia absinthium* and *Artemisia vulgaris*. *Chemistry Central Journal*, *8*(1). <https://doi.org/10.1186/1752-153X-8-6>.
- Pandey, J., Bhusal, S., Nepali, L., Khatri, M., Ramdam, R., Barakoti, H., Giri, P. M., Pant, D., Aryal, P., Rokaya, R. K., & Bhandari, R. (2021). Anti-Inflammatory Activity of *Artemisia vulgaris* Leaves, Originating from Three Different Altitudes of Nepal. *Scientific World Journal*, 2021. <https://doi.org/10.1155/2021/6678059>.
- Pujiastuti, E., & Ma'rifah, S. (2022). Pengaruh Pengeringan Terhadap Kadar Total Flavonoid dan Aktivitas Antioksidan Ekstrak Etanol 70% Daun Jamblang (*Syzygium cumini*). *Jurnal Ilmu Kefarmasian*, *3*(2), 318–324. <https://doi.org/https://doi.org/10.31764/lf.v3i2.9379>.
- Puspita Sari, D., Yuniar, S., Awalia Nur Fadillah, S., Mutiarani, A., & Kusumawaty, D. (2021). The Effectiveness of Mugwort Leaf Extract and Gotu kola Leaf Extract against Acne Bacterial Activity. <https://doi.org/10.17509/xxxx.xxxx>.
- Rahmawati, I., Fachri, B. A., Nurtsulutsiyah, N., Manurung, Y. H., Muhammad, R., Palupi, B., Rizkiana, M. F., & Amini, H. W. (2022). Penerapan Response Surface Methodology dalam Optimasi Kondisi Proses Ekstraksi Antosianin pada Limbah Kulit Kakao dengan Metode Maserasi Menggunakan Pelarut Etanol. *JC-T (Journal Cis-Trans): Jurnal Kimia Dan Terapannya*, *6*(1), 24–31. <https://doi.org/10.17977/um0260v6i12022p024>.
- Reungoat, V., Chadni, M., & Ioannou, I. (2021). Response Surface Methodology Applied to the Optimization of Phenolic Compound Extraction from *Brassica*. In P. Kayaroganam (Ed.), *Response Surface Methodology in Engineering Science*. IntechOpen. <https://doi.org/10.5772/intechopen.97655>.

- Rivai, H., Azmel, S. N. F., & Yetti, R. D. (2020). Analisis Fitokimia Ramuan Obat Tradisional Untuk Nyeri Haid: Baru Cina (*Artemisia vulgaris* (L.)). *Research Gate*. <https://doi.org/10.13140/RG.2.2.16119.44965>.
- Sari, M., Ulfa, R. N., Marpaung, M. P., & Purnama. (2021). Penentuan Aktivitas Antioksidan dan Kandungan Flavonoid Total Ekstrak Daun Papasan (*Coccinia grandis* L.) Berdasarkan Perbedaan Pelarut Polar. *KOVALEN: Jurnal Riset Kimia*, 7(1), 30–41. <https://doi.org/https://doi.org/10.22487/kovalen.2021.v7.i1.15437>.
- Suharyanto, & Prima D.A.N. (2020). Penetapan Kadar Flavonoid Toral pada Juice Daun Ubi Jalar Ungu (*Ipomoea batatas* L.) yang Berpotensi sebagai Hepatoprotektor dengan Metode Spektrofotometri UV-Vis. *Cendekia Journal of Pharmacy*, 2(4), 110–119. <https://doi.org/https://doi.org/10.31596/cjp.v4i2>.
- Syamsul, E. S., Hakim, Y. Y., & Nurhasnawati, H. (2019). Penetapan Kadar Flavonoid Ekstrak Daun Kelakai (*Stenochlaena palustris* (Burm. F.) Bedd.) dengan Metode Spektrofotometri UV-VIS (Vol. 1, Issue 1).
- Tiwari, R. H., Ahmad, A., Khan, A. F., Al-Keridis, L. A., Saeed, M., Alshammari, N., Alabdallah, N. M., Ansari, I. A., & Mujeeb, F. (2023). Ethanolic Extract of *Artemisia vulgaris* Leaf Promotes Apoptotic Cell Death in Non-Small-Cell Lung Carcinoma A549 Cells through Inhibition of the Wnt Signaling Pathway. *Metabolites*, 13(4), 1–19. <https://doi.org/10.3390/metabo13040480>.
- Trifan, A., Czerwińska, M. E., Mardari, C., Zengin, G., Sinan, K. I., Korona-Głowniak, I., Skalicka-Woźniak, K., & Luca, S. V. (2022). Exploring the *Artemisia* Genus: An Insight into the Phytochemical and Multi-Biological Potential of *A. campestris* subsp. *lednicensis* (Spreng.) Greuter & Raab-Straube. *Plants*, 11(21). <https://doi.org/10.3390/plants11212874>.
- Xiao, M., Ye, J., Huang, Y., & Zhang, X. (2014). Response surface optimization for extraction of flavonoids from *Artemisia lactiflora* wall. ex dc. and evaluation of antioxidant capacities in vitro. *Asian Journal of Chemistry*, 26(9), 2802–2808. <https://doi.org/10.14233/ajchem.2014.16962>.